

## ORIGINAL ARTICLE

## Meat intake and risk of diverticulitis among men

Yin Cao,<sup>1,2,3</sup> Lisa L Strate,<sup>4</sup> Brieze R Keeley,<sup>5</sup> Idy Tam,<sup>6</sup> Kana Wu,<sup>3</sup>  
Edward L Giovannucci,<sup>3,7,8</sup> Andrew T Chan<sup>1,2,8,9</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2016-313082>).

For numbered affiliations see end of article.

**Correspondence to**

Dr Andrew T Chan, Clinical and Translational Epidemiology Unit, Massachusetts General Hospital, 55 Fruit Street, Boston, MA 02114, USA; [achan@mgh.harvard.edu](mailto:achan@mgh.harvard.edu)

Received 20 September 2016

Revised 4 November 2016

Accepted 20 November 2016

**ABSTRACT**

**Objective** Diverticulitis is a common disease with a substantial clinical and economic burden. Besides dietary fibre, the role of other foods in the prevention of diverticulitis is underexplored.

**Design** We prospectively examined the association between consumption of meat (total red meat, red unprocessed meat, red processed meat, poultry and fish) with risk of incident diverticulitis among 46 461 men enrolled in the Health Professionals Follow-Up Study (1986–2012). Cox proportional hazards models were used to compute relative risks (RRs) and 95% CIs.

**Results** During 651 970 person-years of follow-up, we documented 764 cases of incident diverticulitis. Compared with men in the lowest quintile (Q1) of total red meat consumption, men in the highest quintile (Q5) had a multivariable RR of 1.58 (95% CI 1.19 to 2.11; *p* for trend=0.01). The increase in risk was non-linear, plateauing after six servings per week (*p* for non-linearity=0.002). The association was stronger for unprocessed red meat (RR for Q5 vs Q1: 1.51; 95% CI 1.12 to 2.03; *p* for trend=0.03) than for processed red meat (RR for Q5 vs Q1: 1.03; 95% CI 0.78 to 1.35; *p* for trend=0.26). Higher consumption of poultry or fish was not associated with risk of diverticulitis. However, the substitution of poultry or fish for one serving of unprocessed red meat per day was associated with a decrease in risk of diverticulitis (multivariable RR 0.80; 95% CI 0.63 to 0.99).

**Conclusions** Red meat intake, particularly unprocessed red meat, was associated with an increased risk of diverticulitis. The findings provide practical dietary guidance for patients at risk of diverticulitis.

**INTRODUCTION**

Diverticulitis is the inflammation of diverticula of the colon. It is a common disease that results in about 210 000 hospitalisations per year in the USA at a cost of more than US\$2 billion.<sup>1</sup> Recently, the incidence of diverticulitis has been rising, particularly in young individuals.<sup>2–3</sup> Approximately 4% of patients with diverticula develop acute or chronic complications including perforation, abscess and fistula.<sup>4</sup> Despite the enormous clinical and economic burden of diverticulitis, little is known about its epidemiology and etiopathogenesis.<sup>5</sup> Although smoking,<sup>6–7</sup> non-steroidal anti-inflammatory drugs (NSAIDs),<sup>8</sup> physical inactivity and obesity<sup>9–11</sup> are identified as risk factors for diverticulitis, dietary factors are less explored.

Dietary fibre is the most studied dietary risk factor for diverticular disease. However, a few studies suggest that red meat consumption may also

**Significance of this study****What is already known on this subject?**

- Diverticulitis is a common disease that results in enormous clinical and economic burden.
- Little is known about its epidemiology and etiopathogenesis.
- Besides dietary fibre, the role of other dietary factors in the prevention of diverticulitis is underexplored.

**What are the new findings?**

- Red meat intake, particularly unprocessed red meat intake, was associated with an increased risk of diverticulitis.
- Substitution of unprocessed red meat per day with poultry or fish may reduce the risk of diverticulitis.

**How might it impact on clinical practice in the foreseeable future?**

- The findings may provide practical dietary guidance for patients at risk of diverticulitis.

be important.<sup>12–13</sup> A recent prospective UK population-based cohort study found that risk of diverticular disease was 31% lower among vegetarians or vegans (relative risk (RR): 0.69; 95% CI 0.55 to 0.86) compared with meat eaters.<sup>13</sup> However, the endpoint of the study was diverticular disease that required hospitalisation, and therefore the results may not be generalisable to patients with more common and mild presentations of diverticulitis.<sup>1</sup> In addition, the specific contribution of red meat, poultry or fish to the observed link was not investigated.

In our prior analysis from a large prospective cohort study, the Health Professionals Follow-Up Study (HPFS), we found that red meat intake, independent of fibre, may be associated with a composite outcome of symptomatic diverticular disease, which included 385 incident cases over 4 years of follow-up.<sup>12</sup> Diverticulitis is distinct in presentation, treatment and pathophysiology from other manifestations of diverticular disease including uncomplicated diverticulosis and diverticular bleeding.<sup>5</sup> Thus, in the present study, we updated this analysis, which allowed us to prospectively examine the association between consumption of meat (total red meat, red unprocessed meat, red processed meat, poultry and fish) with risk of incident diverticulitis in 764 cases over 26 years of follow-up.

**To cite:** Cao Y, Strate LL, Keeley BR, *et al.* Gut Published Online First: [please include Day Month Year] doi:10.1136/gutjnl-2016-313082

## METHODS

### Study population

The HPFS is a large ongoing prospective cohort study of 51 529 US male health professionals aged 40–75 years at enrolment in 1986. Participants have been mailed questionnaires every 2 years since baseline to collect data on demographics, lifestyle factors, medical history and disease outcomes and every 4 years to update dietary intake. The overall follow-up rate is greater than 94%.<sup>14</sup> This study was approved by the institutional review board at the Harvard T.H. Chan School of Public Health.

Men who reported a diagnosis of diverticulosis or its complications (n=253), IBD (n=502) or a cancer of the GI tract (n=2038) at baseline in 1986 were excluded from the analysis (figure 1). In addition, we excluded study participants who reported implausible energy intakes (<800 or >4200 kcal/day) (n=1504) and who did not answer questions regarding intake of unprocessed and processed meat, poultry and fish (n=748). A total of 46 461 men were included in the current analysis.

### Assessment of meat intake

Participants in the HPFS completed semiquantitative food frequency questionnaires (FFQ) in 1986, which was updated every 4 years. In each FFQ, we asked the participants how often, on average, they consumed each food of a standard portion size during the past year. There were nine possible responses, ranging from ‘never or less than once per month’ to ‘six or more times per day’. For unprocessed red meat consumption, the FFQ included questions on ‘beef or lamb as main dish’, ‘pork as main dish’, ‘hamburger’ and ‘beef, pork or lamb as a sandwich or mixed dish’. For processed red meat, there were questions on ‘bacon’, ‘beef or pork hot dogs’, ‘salami, bologna or other processed meat sandwiches’ and ‘other processed red meats, such as sausage, kielbasa, etc’. We derived total red meat consumption by summing consumption of unprocessed and processed red meat. For total poultry consumption, the FFQ included questions on ‘chicken or turkey with or without skin’, ‘chicken or turkey hot dogs’ and ‘chicken or turkey sandwiches’. For total fish intake, consumption of ‘dark meat fish’, ‘canned tuna fish’, ‘breaded fish cakes, pieces or fish sticks’ and ‘other fish’ were added up. The reproducibility and validity of the FFQs in measuring food intake have been previously described.<sup>15 16</sup>

### Ascertainment of diverticulitis cases

The primary endpoint of this study was incident diverticulitis. Beginning in 1990, participants who reported newly diagnosed diverticulitis or diverticulosis on the biennial study questionnaire were sent supplementary questionnaires that ascertained the date of diagnosis, presenting symptoms, diagnostic procedures and treatment. Diverticulitis was defined as abdominal pain attributed to diverticular disease and one of the following criteria: (1) complicated by perforation, abscess, fistula or obstruction; (2) requiring hospitalisation, antibiotics or surgery or (3) pain categorised as severe or acute or abdominal pain presenting with fever, requiring medication or evaluated using abdominal CT. We have previously used these case definitions and documented the validity of self-reported diverticulitis in this population.<sup>10 11 17</sup>


Beginning in 2006, we administered a revised supplementary diverticular disease questionnaire. Using questions that included definitions for each disease outcome, we assessed uncomplicated diverticulitis, complications of diverticulitis including abscess, fistula, perforation and obstruction, diverticular bleeding and diverticulosis.<sup>8</sup>

### Statistical analysis

In our primary analyses, we examined the association between consumption of meat (total red meat, red unprocessed meat, red processed meat, poultry and fish) with risk of incident diverticulitis. Person-years of follow-up accrued from the date of return of the 1986 questionnaire until either the date of diagnosis of diverticulitis, diverticulosis or diverticular bleeding, death or 31 December 2012, whichever came first. We censored men who reported a new diagnosis of GI cancer or IBD.

Cox proportional hazards models with time-varying meat consumption and covariates were used to compute HRs as estimates for age-adjusted and multivariable-adjusted RRs and 95% CIs. To control as finely as possible for confounding by age, calendar time and a possible interaction between these two time scales, we stratified models jointly by age (in months) and 2-year questionnaire cycle. In age-adjusted models, we additionally adjusted for total energy intake. In multivariable models, we further adjusted for the following potential confounders: body mass index (BMI) (kg/m<sup>2</sup> in quintiles), vigorous physical activity including activities with a metabolic equivalent task (MET) score of 6 or more (MET hours/week in quintiles), smoking history (never smokers, <4.9, 5–19.9, 20–39.9, ≥40 pack

**Figure 1** Flow diagram of the study population.



years), fibre intake (g/day in quintiles), regular use of aspirin, non-aspirin NSAIDs and acetaminophen ( $\geq 2$  vs  $< 2$  times/week). We updated meat consumption as well as other covariates prior to each 2-year or 4-year interval (simple updating).

Tests for linear trend were performed using meat consumption as a continuous variable. We examined the possible non-linear relation between meat consumption and risk of diverticulitis non-parametrically with restricted cubic splines.<sup>18–20</sup> To test for non-linearity, we used a likelihood ratio test, comparing the model with only the linear term to the model with the linear and the cubic spline terms. Departures from the proportional hazards assumption were tested by likelihood ratio tests comparing models with and without the interaction terms of age or calendar time by categories of meat consumption. No significant violation of the proportionality assumption was found ( $p > 0.05$  for all tests).

As an exploratory analysis, we assessed the association between components of red meat including total fat, saturated fat, cholesterol and haem iron and incident diverticulitis, with and without additionally adjusting for unprocessed and processed red meat. To facilitate the translation to dietary recommendations regarding meat intake, we estimated the associations of substituting one serving of poultry or fish for one serving of red meat with incident diverticulitis by including both as continuous variables in the same multivariate model. The difference in their  $\beta$  coefficients, as well as their variances and covariance, was used to estimate the RR and 95% CI for the substitution associations.<sup>21 22</sup> All of the analyses were performed using SAS V.9.4 (SAS Institute, Cary, North Carolina, USA), and the statistical tests were two-sided and  $p$  values less than 0.05 were considered statistically significant.

## RESULTS

During 651 970 person-years of follow-up, we documented 764 incident cases of diverticulitis.

Compared with men with lower intake of red meat, men with higher red meat consumption smoked more, used non-aspirin NSAIDs and acetaminophen more often, and were less likely to

exercise vigorously (table 1). As expected, their intake of total fat, saturated fat, cholesterol and haem iron were substantially higher. In contrast, fibre intake among these men was lower. Compared with men with lower poultry or /fish consumption, men with higher poultry or /fish consumption were more likely to be engaged in vigorous physical activity, use aspirin, smoke less, and had higher intake of haem iron. Men with higher poultry intake also had higher intake of cholesterol, and individuals with higher fish intake had lower intake of total and saturated fat.

Overall, total red meat intake was associated with an increased risk of diverticulitis. Compared with men in the lowest quintile (Q1) of total red meat consumption, men in the highest quintile (Q5) had a multivariable RR of 1.58 (95% CI 1.19 to 2.11) (table 2) after adjustment for total fibre and all other potential confounding variables. The risk of incident diverticulitis increased 18% per serving of red meat consumed per day ( $p$  for trend=0.01). Nonparametric regression curves suggest that the dose–response relationship for total red meat was non-linear ( $p$  for non-linearity=0.002): even one serving per week appeared to increase risk, with risk plateauing after six servings per week (figure 2).

The observed link between total red meat intake and risk of diverticulitis appeared primarily driven by consumption of unprocessed red meat. Men in the Q5 of unprocessed meat consumption had an RR of 1.51 (95% CI 1.12 to 2.03;  $p$  for trend=0.03) compared with men in the Q1, even after adjusting for processed red meat intake. In contrast, processed red meat was associated with increased risk of diverticulitis only in the age-adjusted model, but not after further controlling for other covariates, without or with adjustment for unprocessed red meat (RR for the Q5 vs Q1: 1.03; 95% CI 0.78 to 1.35;  $p$  for trend=0.26). Additional adjustment for red meat components including total fat, saturated fat, cholesterol and haem iron minimally changed the associations observed above (data not shown). The findings were similar among overweight and obese men, and men who were younger than 60 years or aged 60 years and above.

**Table 1** Characteristics according to person-years of meat intake consumption, Health Professionals Follow-Up Study (HPFS) 1986–201.2

	Meat intake, quintile														
	Total red meat			Unprocessed meat			Processed meat			Poultry			Fish		
	1	3	5	1	3	5	1	3	5	1	3	5	1	3	5
Meat intake, servings/week	1.2	5.3	13.5	0.8	3.2	8.6	0	1.4	6.4	0.7	2.7	6.2	0.3	1.6	2.5
Age, years*	63	62	62	64	62	61	63	62	62	65	62	60	62	61	63
BMI, kg/m <sup>2</sup>	25	26	27	25	26	26	25	26	26	26	26	26	26	26	26
Vigorous physical activity, MET-hours/week	18	12	9	18	12	9	17	12	9	12	13	14	10	12	14
Ever smokers, %	46	51	55	47	52	52	44	51	56	51	51	49	49	51	51
Pack-year among ever smokers	21	24	28	22	25	27	21	24	28	26	24	23	27	25	23
Alcohol intake, g/day	8.8	12	13	9	12	13	8.7	12	14	11	12	11	10	12	12
Dietary fibre, g/day	28	22	20	27	22	20	28	22	20	23	23	23	21	22	24
Total fat, g/day	58	70	78	60	70	77	59	70	77	70	68	68	72	70	68
Saturated fat, g/day	17	23	27	18	23	26	18	23	26	23	22	21	25	23	22
Cholesterol, mg/day	213	267	320	219	268	313	217	266	312	238	263	293	257	269	266
Haem iron, mg/day	0.8	1.2	1.5	0.8	1.1	1.6	0.9	1.2	1.3	1.0	1.2	1.4	1.1	1.2	1.2
Regular aspirin use, %	47	46	45	47	47	46	47	47	45	44	47	48	44	45	49
Regular non-aspirin NSAID use, %	12	15	16	12	15	16	12	15	15	13	14	15	15	14	14
Regular acetaminophen use, %	6.6	8.3	9.4	6.8	8.2	9.3	6.4	8.1	9.2	7.3	7.9	8.8	8.5	8.1	8.3

\*All values other than age have been directly standardised to age distribution (in 5-year age group) of all the participants. Mean was presented for continuous variables. MET, metabolic equivalent task.

**Table 2** Meat consumption and risk of incident diverticulitis

	Meat consumption, quintile					Per 1 serving/day	p For trend*
	1	2	3	4	5		
<b>Total red meat</b>							
Cases, No.	106	158	164	153	183		
Person-years	134 819	128 596	128 853	127 720	131 982		
Median intake, servings/week	1.5	3.4	5.4	7.9	12.4		
Age-adjusted RR (95% CI)†	1 (Reference)	1.56 (1.22 to 2.01)	1.69 (1.32 to 2.18)	1.67 (1.29 to 2.16)	2.09 (1.60 to 2.71)	1.33 (1.19 to 1.48)	<0.001
Multivariable RR (95% CI)‡	1 (Reference)	1.39 (1.08 to 1.80)	1.43 (1.10 to 1.85)	1.35 (1.02 to 1.77)	1.58 (1.19 to 2.11)	1.18 (1.04 to 1.33)	0.01
<b>Unprocessed red meat</b>							
Cases, No.	99	148	189	162	166		
Person-years	135 137	121 492	136 654	126 194	132 493		
Median intake, servings/week	0.98	2.0	2.9	5.0	8.0		
Age-adjusted RR (95% CI)†	1 (Reference)	1.63 (1.26 to 2.11)	1.90 (1.48 to 2.44)	1.85 (1.42 to 2.39)	1.95 (1.49 to 2.55)	1.48 (1.26 to 1.75)	<0.001
Multivariable RR (95% CI)‡	1 (Reference)	1.46 (1.12 to 1.90)	1.63 (1.26 to 2.11)	1.52 (1.16 to 1.99)	1.54 (1.16 to 2.05)	1.26 (1.05 to 1.51)	0.01
Multivariable RR (95% CI)§	1 (Reference)	1.47 (1.12 to 1.93)	1.61 (1.23 to 2.10)	1.49 (1.13 to 1.98)	1.51 (1.12 to 2.03)	1.23 (1.02 to 1.49)	0.03
<b>Processed red meat</b>							
Cases, No.	116	130	184	165	169		
Person-years	120 357	127 492	155 568	118 180	130 373		
Median intake, servings/week	0	0.5	1.5	2.5	5.5		
Age-adjusted RR (95% CI)†	1 (Reference)	1.12 (0.87 to 1.45)	1.26 (1.00 to 1.60)	1.54 (1.21 to 1.97)	1.49 (1.16 to 1.91)	1.35 (1.14 to 1.58)	<0.001
Multivariable RR (95% CI)‡	1 (Reference)	1.00 (0.77 to 1.30)	1.05 (0.83 to 1.35)	1.23 (0.95 to 1.59)	1.13 (0.87 to 1.48)	1.15 (0.96 to 1.38)	0.14
Multivariable RR (95% CI)§	1 (Reference)	0.95 (0.73 to 1.23)	0.95 (0.74 to 1.23)	1.10 (0.84 to 1.43)	1.03 (0.78 to 1.35)	1.11 (0.92 to 1.35)	0.26
<b>Poultry</b>							
Cases, No.	145	160	186	127	146		
Person-years	130 083	137 482	129 756	115 726	138 924		
Median intake, servings/week	1.0	1.5	3.0	3.5	6.0		
Age-adjusted RR (95% CI)†	1 (Reference)	1.15 (0.91 to 1.45)	1.30 (1.04 to 1.63)	1.08 (0.84 to 1.38)	1.05 (0.83 to 1.34)	1.03 (0.82 to 1.28)	0.83
Multivariable RR (95% CI)‡	1 (Reference)	1.13 (0.89 to 1.43)	1.33 (1.06 to 1.66)	1.13 (0.88 to 1.45)	1.09 (0.86 to 1.39)	1.07 (0.86 to 1.33)	0.55
<b>Fish</b>							
Cases, No.	159	152	173	141	139		
Person-years	117 176	113 430	163 700	122 327	135 337		
Median intake, servings/week	0.5	1.0	1.5	2.5	4.5		
Age-adjusted RR (95% CI)†	1 (Reference)	1.01 (0.80 to 1.26)	0.82 (0.66 to 1.03)	0.83 (0.66 to 1.05)	0.76 (0.60 to 0.97)	0.69 (0.51 to 0.93)	0.01
Multivariable RR (95% CI)‡	1 (Reference)	1.03 (0.82 to 1.29)	0.86 (0.69 to 1.08)	0.90 (0.71 to 1.14)	0.87 (0.68 to 1.10)	0.82 (0.61 to 1.11)	0.20

\*Calculated using continuous meat intake.

†Adjusted for age, questionnaire cycle and total energy intake (quintiles).

‡Additionally adjusted for BMI (quintiles), vigorous physical activity (quintiles), smoking history (never smokers, &lt;4.9, 5–19.9, 20–39.9, ≥40 pack years), fibre intake (quintiles), regular use of aspirin, non-aspirin NSAIDs and acetaminophen.

§Additionally adjusted for processed red meat intake or unprocessed red meat intake (quintiles).

An exploratory analysis of major red meat components indicated a modest association between total fat intake and diverticulitis (RR for the Q5 vs Q1: 1.33; 95% CI 1.00 to 1.77) after adjusting for unprocessed and processed red meat intake (see online supplementary table S1). Poultry was not associated with risk of incident diverticulitis (RR for the Q5 vs Q1: 1.09; 95% CI 0.86 to 1.39; *p* for trend=0.55) (table 2). Higher fish intake was associated with reduced risk of diverticulitis in age-adjusted model, but not after further adjustment for other potential confounders (RR for the Q5 vs Q1: 0.87; 95% CI 0.68 to 1.10; *p* for trend=0.20). The multivariable RR of diverticulitis associated with substitution of poultry or fish for one serving of unprocessed red meat per day was 0.80 (95% CI 0.63 to 0.99). In contrast, substituting one serving of red processed meat per day with poultry or fish was not significantly associated with risk of diverticulitis (RR 0.86; 95% CI 0.67 to 1.09).


## DISCUSSION

In this large prospective cohort of men, total red meat intake, especially consumption of unprocessed red meat, was non-linearly associated with an increased risk of diverticulitis.

Moreover, we identified unprocessed red meat, but not processed red meat, as the major driver for the link between red meat and diverticulitis. The association was independent of fibre intake. As a component of red meat, total fat intake was associated with an increased risk of diverticulitis even after adjusting for unprocessed and processed red meat intake. In contrast, higher consumption of poultry or fish was not linked with risk of incident diverticulitis. However, substitution of one serving of unprocessed red meat per day with poultry or fish was associated with a 20% lower risk of diverticulitis.

Our findings were generally in line with our prior early analysis that red meat, but not poultry or fish, is associated with an increased risk of diverticular disease. However, in this study we were able to examine diverticulitis separately from diverticular bleeding and symptomatic uncomplicated diverticular disease, rather than using a composite endpoint. In addition, in comparison with the prior study of red meat in the HPFS cohort,<sup>12</sup> our cohort included 22 additional years of follow-up and more than twice as many cases.

Pathways through which red meat consumption may influence risk of diverticulitis are yet to be established. Chronic low-grade



**Figure 2** Non-parametric restricted cubic splines of meat intake (servings/week) and risk of diverticulitis. (A) Total red meat. (B) Unprocessed red meat. (C) Processed red meat. (D) Poultry. (E) Fish.

systemic inflammation may be an essential step.<sup>5</sup> Higher red meat intake is associated with higher levels of inflammatory biomarkers such as C-reactive protein and ferritin,<sup>23</sup> as well as an increased risk of chronic diseases such as cardiovascular disease, diabetes and cancer, in which chronic inflammation has been implicated in the pathogenesis.<sup>24–26</sup> The gut microbiome may also mediate the link between red meat and diverticulitis. Emerging evidence suggest that short-term and long-term diet, particularly red meat intake, alters microbial community structure, overwhelms interindividual differences in microbial gene expression and changes the metabolism of bacteria in the colon.<sup>27</sup> Although a direct link between the gut microbiome and diverticulitis is yet to be established,<sup>28</sup> it is recently hypothesised that changes in intestinal microbiota composition may play a similar role in the development of diverticular disease and its complications.<sup>29</sup> Specifically, changes in the microbiota composition lead to deficiencies of host immune defences and dysfunction of the mucosal barrier resulting in increased mucosal adherence and translocation of bacteria. A pathogenic immune response is then activated and the release of proinflammatory cytokines further induces inflammation, leading to symptoms.<sup>30</sup>

We also observed that unprocessed red meat, but not processed red meat, was the primary driver for the association between total red meat and risk of diverticulitis. Compared with processed meat, unprocessed meat (eg, steak) is usually consumed in larger portions, which could lead to a larger undigested piece in the large bowel and induce different changes in colonic microbiota. In addition, higher cooking temperatures used in the preparation of unprocessed meat may influence bacterial composition or proinflammatory mediators in the colon. These hypotheses need to be confirmed by other studies.

The strengths of our study include a large, well-characterised population with detailed, prospective and updated assessment of meat consumption over 26 years of follow-up. We were also able to differentiate diverticulitis from diverticular bleeding and uncomplicated diverticulosis. As these manifestations appear to arise via different biological mechanisms, they are likely to have distinct risk factors. In addition, the large number of cases of diverticulitis accrued during long-term follow-up in this study allowed us to examine subtype-specific meat consumption and its dose–response relationship with risk of incident diverticulitis.

There are also limitations of this study. First, misclassification of self-reported outcome was likely. However, healthcare professionals are more likely to accurately self-report medical information, and reports of diverticulitis have been validated in this cohort.<sup>10 11 17</sup> Second, measurement errors associated with recall of meat consumption as well as potential confounders were possible; however, they would be non-differential to diagnosis of diverticulitis. Third, even though we were able to adjust for a variety of potential confounders, the possibility of residual confounding can never be ruled out. In addition, due to limited number of people consuming a vegetarian diet, we were unable to estimate the substitution effect of a vegetarian dish. Finally, the generalisability of our data to other populations, particularly women and other racial or ethnic groups, may be limited.

## CONCLUSION

In summary, we found that intake of red meat, particularly unprocessed red meat, was associated with an increased risk of diverticulitis. Substitution of unprocessed red meat with poultry or fish may reduce the risk of diverticulitis. Our findings may provide practical dietary guidance for patients at risk of diverticulitis, a common disease of huge economic and clinical

burden. The mechanisms underlying the observed associations require further investigation.

## Author affiliations

<sup>1</sup>Clinical and Translational Epidemiology Unit, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA

<sup>2</sup>Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA

<sup>3</sup>Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

<sup>4</sup>Division of Gastroenterology, University of Washington School of Medicine, Harborview Medical Center, Seattle, Washington, USA

<sup>5</sup>Department of Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA

<sup>6</sup>Tufts University School of Medicine, Boston, Massachusetts, USA

<sup>7</sup>Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

<sup>8</sup>Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

<sup>9</sup>Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA

**Acknowledgements** We would like to thank the participants and staff of the Health Professionals Follow-Up Study for their valuable contributions.

**Contributors** YC and ATC had full access to all of the data in the study, and take responsibility for the integrity of the data and the accuracy of the data analysis. Study concept and design: YC, LLS, ELG and ATC. Acquisition of data: LLS, BRK, IT and ATC. Analysis and interpretation of data: all coauthors. Drafting of the manuscript and statistical analysis: YC. Critical revision of the manuscript for important intellectual content: all coauthors. Obtained funding: LLS, ELG and ATC. Administrative, technical or material support and study supervision: LLS and ATC.

**Funding** This work was supported by grants R01 DK101495, R01 DK084157, K24 DK098311 and UM1 CA167552 from the National Institutes of Health.

**Competing interests** ATC previously served as a consultant for Bayer Healthcare, Aralaz Pharmaceuticals and Pfizer Inc. for work unrelated to the topic of this manuscript. This study was not funded by Bayer Healthcare, Pozen Aralaz Pharmaceuticals or Pfizer Inc.

**Ethics approval** Harvard T.H. Chan School of Public Health.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- 1 Peery AF, Crockett SD, Barritt AS, *et al.* Burden of gastrointestinal, liver, and pancreatic diseases in the United States. *Gastroenterology* 2015;149:1731–1741. e3.
- 2 Nguyen GC, Sam J, Anand N. Epidemiological trends and geographic variation in hospital admissions for diverticulitis in the United States. *World J Gastroenterol* 2011;17:1600–5.
- 3 Bharucha AE, Parthasarathy G, Ditha I, *et al.* Temporal trends in the incidence and natural history of diverticulitis: a population-based study. *Am J Gastroenterol* 2015;110:1589–96.
- 4 Shahedi K, Fuller G, Bolus R, *et al.* Long-term risk of acute diverticulitis among patients with incidental diverticulosis found during colonoscopy. *Clin Gastroenterol H* 2013;11:1609–13.
- 5 Strate LL, Modi R, Cohen E, *et al.* Diverticular disease as a chronic illness: evolving epidemiologic and clinical insights. *Am J Gastroenterol* 2012;107:1486–93.
- 6 Hjern F, Wolk A, Håkansson N. Smoking and the risk of diverticular disease in women. *Br J Surg* 2011;98:997–1002.
- 7 Humes DJ, Fleming KM, Spiller RC, *et al.* Concurrent drug use and the risk of perforated colonic diverticular disease: a population-based case-control study. *Gut* 2011;60:219–24.
- 8 Strate LL, Liu YL, Huang ES, *et al.* Use of aspirin or nonsteroidal anti-inflammatory drugs increases risk for diverticulitis and diverticular bleeding. *Gastroenterology* 2011;140:1427–33.
- 9 Hjern F, Wolk A, Håkansson N. Obesity, physical inactivity, and colonic diverticular disease requiring hospitalization in women: a prospective cohort study. *Am J Gastroenterol* 2012;107:296–302.
- 10 Strate LL, Liu YL, Aldoori WH, *et al.* Physical activity decreases diverticular complications. *Am J Gastroenterol* 2009;104:1221–30.
- 11 Strate LL, Liu YL, Aldoori WH, *et al.* Obesity increases the risks of diverticulitis and diverticular bleeding. *Gastroenterology* 2009;136:115–22.
- 12 Aldoori WH, Giovannucci EL, Rimm EB, *et al.* A prospective study of diet and the risk of symptomatic diverticular-disease in men. *Am J Clin Nutr* 1994;60:757–64.
- 13 Crowe FL, Appleby PN, Allen NE, *et al.* Diet and risk of diverticular disease in Oxford cohort of European Prospective Investigation into Cancer and Nutrition (EPIC): prospective study of British vegetarians and non-vegetarians. *BMJ* 2011;343.

- 14 Rimm EB, Stampfer MJ, Colditz GA, *et al.* Effectiveness of various mailing strategies among nonrespondents in a prospective cohort study. *Am J Epidemiol* 1990;131:1068–71.
- 15 Hu FB, Rimm E, Smith-Warner SA, *et al.* Reproducibility and validity of dietary patterns assessed with a food-frequency questionnaire. *Am J Clin Nutr* 1999;69:243–9.
- 16 Feskanich D, Rimm EB, Giovannucci EL, *et al.* Reproducibility and validity of food intake measurements from a semiquantitative food frequency questionnaire. *J Am Diet Assoc* 1993;93:790–6.
- 17 Strate LL, Liu YL, Syngal S, *et al.* Nut, corn, and popcorn consumption and the incidence of diverticular disease. *JAMA* 2008;300:907–14.
- 18 Smith PL. Splines as a Useful and Convenient Statistical Tool. *Am Stat* 1979;33:57–62.
- 19 Harrell FE Jr, Lee KL, Pollock BG. Regression models in clinical studies: determining relationships between predictors and response. *J Natl Cancer Inst* 1988;80:1198–202.
- 20 Durrleman S, Simon R. Flexible regression-models with cubic-splines. *Stat Med* 1989;8:551–61.
- 21 Halton TL, Willett WC, Liu S, *et al.* Potato and French fry consumption and risk of type 2 diabetes in women. *Am J Clin Nutr* 2006;83:284–90.
- 22 Bernstein AM, Sun Q, Hu FB, *et al.* Major dietary protein sources and risk of coronary heart disease in women. *Circulation* 2010;122:876–83.
- 23 Ley SH, Sun Q, Willett WC, *et al.* Associations between red meat intake and biomarkers of inflammation and glucose metabolism in women. *Am J Clin Nutr* 2014;99:352–60.
- 24 Micha R, Michas G, Mozaffarian D. Unprocessed red and processed meats and risk of coronary artery disease and type 2 diabetes—an updated review of the evidence. *Curr Atheroscler Rep* 2012;14:515–24.
- 25 Pan A, Sun Q, Bernstein AM, *et al.* Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *Am J Clin Nutr* 2011;94:1088–96.
- 26 Chan DS, Lau R, Aune D, *et al.* Red and processed meat and colorectal cancer incidence: meta-analysis of prospective studies. *PLoS ONE* 2011;6:e20456.
- 27 David LA, Maurice CF, Carmody RN, *et al.* Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 2014;505:559–63.
- 28 Gueimonde M, Ouwehand A, Huhtinen H, *et al.* Qualitative and quantitative analyses of the bifidobacterial microbiota in the colonic mucosa of patients with colorectal cancer, diverticulitis and inflammatory bowel disease. *World J Gastroenterol* 2007;13:3985–9.
- 29 Daniels L, Philipszoon LE, Boermeester MA. A hypothesis: important role for gut microbiota in the etiopathogenesis of diverticular disease. *Dis Colon Rectum* 2014;57:539–43.
- 30 Guarner F, Malagelada JR. Gut flora in health and disease. *Lancet* 2003;361:512–19.



## Meat intake and risk of diverticulitis among men

Yin Cao, Lisa L Strate, Brieze R Keeley, Idy Tam, Kana Wu, Edward L Giovannucci and Andrew T Chan

*Gut* published online January 9, 2017

---

Updated information and services can be found at:

<http://gut.bmj.com/content/early/2017/01/03/gutjnl-2016-313082>

---

*These include:*

### Supplementary Material

Supplementary material can be found at:

<http://gut.bmj.com/content/suppl/2017/01/10/gutjnl-2016-313082.DC1.html>

### References

This article cites 29 articles, 9 of which you can access for free at:

<http://gut.bmj.com/content/early/2017/01/03/gutjnl-2016-313082#BIBL>

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Topic Collections

Articles on similar topics can be found in the following collections

[Press releases](#) (38)

---

### Notes

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>